



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\sin IHL = \frac{1}{3}\sqrt{3}, \cos IHL = \frac{1}{3}\sqrt{6}.$$

$$\sin EHL = \frac{1}{3}, \cos EHL = \frac{2}{3}\sqrt{2}.$$

$$HI : HL = \sin ILH : \sin HIL = \sin EHL : \sin IHE. \therefore HI : w = \frac{1}{3} : \frac{1}{3}\sqrt{6}$$

or $HI = w/\sqrt{6} = w\sqrt{6}/6$; $HE : w = \sin IHL : \sin HIL$; $HE : w = \frac{1}{3}\sqrt{3} : \frac{1}{3}\sqrt{6}$, or
 $HE = w/\sqrt{2} = w\sqrt{2}/2$.

Also solved by G. W. Greenwood.

AVERAGE AND PROBABILITY.


163. Proposed by R. D. CARMICHAEL, Anniston, Ala.

In a regular n -gon a triangle is formed by taking three vertices at random. What is the mean value of the triangle?


Solution by G. B. M. ZERR, Ph. D., Parsons, W. Va.

Whether n be even or odd, there can be formed, at any vertex, as B , or C , or D , etc., by joining it with A and any other vertex, $(n-2)$ triangles. Since there are n vertices, the total number of triangles $= n(n-2)$.

I. n even. For A the sum of the areas is the area of the polygon $= \frac{1}{2}nr^2 \sin(2\pi/n)$; for B and J combined $nr^2 \sin(2\pi/n) =$ sum of areas; for C and I , the sum is $2nr^2 \sin(2\pi/n) - 4(n-2)r^2 \sin(\pi/n) \sin(\pi/n) \sin(2\pi/n)$; for D and H , $3nr^2 \sin(2\pi/n) - 4nr^2 \sin(\pi/n) \sin(\pi/n) \sin(2\pi/n) - 4(n-4)r^2 \sin(\pi/n) \times \sin(2\pi/n) \sin(3\pi/n)$. For the next pair of vertices the sum is



n even.



n odd.

$$\begin{aligned} &4n^2 \sin(2\pi/n) - 4(n+2)r^2 \sin(\pi/n) \sin(\pi/n) \sin(2\pi/n) \\ &- 4(n-2)r^2 \sin(\pi/n) \sin(2\pi/n) \sin(3\pi/n) \\ &- 4(n-6)r^2 \sin(\pi/n) \sin(3\pi/n) \sin(4\pi/n). \end{aligned}$$

The sum of the areas of the triangles for $\frac{n-2}{2}$ th, the $\frac{n}{2}$ th and the $\frac{n+2}{2}$ th vertices are equal, and their combined sum for the three vertices is

$$\begin{aligned}
& \frac{3(n-2)}{4} r^2 \sin(2\pi/n) - 6(2n-8) r^2 \sin(\pi/n) \sin(\pi/n) \sin(2\pi/n) \\
& - 6(2n-12) r^2 \sin(\pi/n) \sin(2\pi/n) \sin(3\pi/n) \\
& - 6(2n-16) r^2 \sin(\pi/n) \sin(3\pi/n) \sin(4\pi/n) \\
& - \dots - 6(4) r^2 \sin(\pi/n) \sin \frac{n-4}{2n} \pi \sin \frac{n-2}{2n} \pi.
\end{aligned}$$

The total sum for all is

$$\begin{aligned}
& \left[\frac{1}{2}n + n(1+2+3+\dots + \frac{n-2}{2}) + \frac{n(n-2)}{4} \right] r^2 \sin(2\pi/n) \\
& - [4r^2(n-2+n+n+2+n+4+\dots + 2n-8) \\
& + 2r^2(2n-8)] \sin(\pi/n) \sin(\pi/n) \sin(2\pi/n) \\
& - [4r^2(n-4+n-2+n+n+2+\dots + 2n-12) \\
& + 2r^2(2n-12)] \sin(\pi/n) \sin(2\pi/n) \sin(3\pi/n) - \text{etc.}, \\
& = \frac{1}{8} n^3 r^2 \sin(2\pi/n) - 3r^2 \sin(\pi/n) [(n-2)(n-4) \sin(\pi/n) \sin(2\pi/n) \\
& + (n-4)(n-6) \sin(2\pi/n) \sin(3\pi/n) + (n-6)(n-8) \sin(3\pi/n) \sin(4\pi/n) + \dots \\
& + 8 \sin \frac{n-4}{2n} \pi \sin \frac{n-2}{2n} \pi] \\
& = \frac{1}{8} n^3 r^2 \sin(2\pi/n) - 24r^2 [\cos(\pi/n) \cos(2\pi/n) + 3\cos(2\pi/n) \cos(3\pi/n) \\
& + 6\cos(3\pi/n) \cos(4\pi/n) + 10\cos(4\pi/n) \cos(5\pi/n) + \dots \\
& + \frac{(n-2)(n-4)}{8} \cos \frac{n-4}{2n} \pi \cos \frac{n-2}{2n} \pi] \sin(\pi/n) \\
& = \frac{1}{8} n^3 r^2 \sin(2\pi/n) - 12r^2 \sin(\pi/n) \cos(\pi/n) (1+3+6+10+\dots + \frac{(n-2)(n-4)}{8}) \\
& - 12r^2 \sin(\pi/n) [\cos(3\pi/n) + 3\cos(5\pi/n) + 6\cos(7\pi/n) + \dots \\
& + \frac{(n-2)(n-4)}{8} \cos \frac{n-3}{n} \pi] = \frac{1}{8} n^3 r^2 \sin(2\pi/n) - \frac{1}{8} n(n-2)(n-4) r^2 \sin(2\pi/n) \\
& + \frac{3n[(n-2) \cos(2\pi/n) - (n-4)] r^2 \sin(2\pi/n)}{8[\sin(\pi/n)]^2} \\
& = \frac{r^2 n \{ (3n-2) [\sin(\pi/n)]^2 - 1 \} \cos(\pi/n)}{2\sin(\pi/n)} \\
\therefore \text{Average area is } & \frac{r^2 \cos(\pi/n)}{2(n-2) \sin(\pi/n)} \{ [3n-2] [\sin(\pi/n)]^2 - 1 \}.
\end{aligned}$$

II. n odd. By a similar process we easily get for the total sum

$$\begin{aligned}
 & \left[\frac{1}{2}nr^2 + nr^2(1+2+3+4+\dots+\frac{n-1}{2}) \right] \sin(2\pi/n) \\
 & - 4r^2(n-2+n+n+2+n+4+\dots+2n-7) \sin(\pi/n) \sin(\pi/n) \sin(2\pi/n) \\
 & - 4r^2(n-4+n-2+n+n+2+\dots+2n-11) \sin(\pi/n) \sin(2\pi/n) \sin(3\pi/n) - \text{etc.}, \\
 & = \frac{n^3+3n}{8} r^2 \sin(2\pi/n) - 3r^2 \sin(\pi/n) [(n-1)(n-3) \sin(\pi/n) \sin(2\pi/n) \\
 & + (n-3)(n-5) \sin(2\pi/n) \sin(3\pi/n) + (n-5)(n-7) \sin(3\pi/n) \sin(5\pi/n) + \dots \\
 & + 8 \sin \frac{n-3}{2n} \pi \sin \frac{n-1}{2n} \pi] \\
 & = \frac{n^3+3n}{8} r^2 \sin(2\pi/n) - 24r^2 \sin(\pi/n) [\cos(\pi/2n) \cos(3\pi/n) \\
 & + 3\cos(3\pi/n) \cos(5\pi/n) + 6\cos(5\pi/n) \cos(7\pi/n) + \dots \\
 & + \frac{(n-1)(n-3)}{8} \cos \frac{n-4}{2n} \pi \cos \frac{n-2}{2n} \pi] = \frac{n^3+3n}{8} r^2 \sin(2\pi/n) \\
 & - 12r^2 \sin(\pi/n) \cos(\pi/n) [1+3+6+10+\dots+\frac{(n-1)(n-3)}{8}] \\
 & - 12r^2 \sin(\pi/n) [\cos(2\pi/n) + 3\cos(4\pi/n) + 6\cos(6\pi/n) + \dots \\
 & + \frac{(n-1)(n-3)}{8} \cos \frac{(n-3)\pi}{n}] = \frac{n^3+3n}{8} r^2 \sin(2\pi/n) \\
 & - \frac{1}{8}(n+1)(n-1)(n-3)r^2 \sin(2\pi/n) + \frac{3r^2}{2\sin(\pi/n)} \\
 & - \frac{[3(n^3-2n-3)\sin(2\pi/n) - 3(n^2-1)\sin(2\pi/n)\cos(2\pi/n)]r^2}{8[\sin(\pi/n)]^2} \\
 & = \frac{r^2}{4\sin(\pi/n)} \{ (4n+3-3n^2) [\sin(\pi/n)]^2 \cos(\pi/n) + 6+6(n+1)\cos(\pi/n) \}.
 \end{aligned}$$

The average area is

$$\frac{r^2}{4n(n-2)\sin(\pi/n)} \{ 6+6(n+1)\cos(\pi/n) - (3n^2-4n-3)\cos(\pi/n) [\sin(\pi/n)]^2 \}.$$

For the foregoing solutions the following explanation is necessary.

$$2\sin \frac{(p-q)\pi}{n} \sin \frac{(p-q+1)\pi}{n} = \cos(\pi/n) + \cos \frac{(2p-2q+1)\pi}{n}$$

$$2\cos \frac{(p-q)\pi}{2n} \cos \frac{(p-q+2)\pi}{2n} = \cos(\pi/n) + \cos \frac{(p-q+1)\pi}{n}$$

$$\text{Let } C = \cos 3\theta + 3\cos 5\theta + 6\cos 7\theta + \dots + \frac{(n-2)(n-4)}{8} \cos(n-3)\theta,$$

$$S = i\sin 3\theta + 3i\sin 5\theta + 6i\sin 7\theta + \dots + \frac{(n-2)(n-4)}{8} i\sin(n-3)\theta,$$

where $\theta = \pi/n$, $i = \sqrt{-1}$.

$$\begin{aligned} \therefore C + S &= \cos 3\theta + i\sin 3\theta + 3(\cos 5\theta + i\sin 5\theta) + 6(\cos 7\theta + i\sin 7\theta) + \dots \\ &+ \frac{(n-2)(n-4)}{8} [\cos(n-3)\theta + i\sin(n-3)\theta] = (\cos\theta + i\sin\theta)^3 + 3(\cos\theta + i\sin\theta)^5 \\ &+ 6(\cos\theta + i\sin\theta)^7 + \dots + \frac{(n-2)(n-4)}{8} (\cos\theta + i\sin\theta)^{n-3} = e^{3i\theta} + 3e^{5i\theta} + 6e^{7i\theta} + \dots \\ &+ \frac{(n-2)(n-4)}{8} e^{(n-3)i\theta} = y^3 [1 + 3y^2 + 6y^4 + 10y^6 + \dots + \frac{(n-2)(n-4)}{8} y^{n-6}] \\ &= \frac{y^3}{(1-y^2)^3} - \frac{(n-1)(n-4)y^{n+3} + n(n-2)y^{n-1} - 2n(n-4)y^{n+1}}{8(1-y^2)^3} \end{aligned}$$

Putting $y = e^{i\theta} = \cos\theta + i\sin\theta$, and equating C = the rational part, we get

$$C = \frac{2n(n-4)\sin(2\pi/n) - n(n-2)\sin(4\pi/n)}{64[\sin(\pi/n)]^3}.$$

MISCELLANEOUS.

164. Proposed by W. J. GREENSTREET, M. A., Editor of The Mathematical Gazette, Stroud, England.

Find the number of real roots of the equation $100\sin x = x$, and show the largest root is approximately 96.10. Find $\tan 39^\circ$ to three places of decimals. How many real roots of $\tan x = 1/x^2$ lie between 0 and 2π ?

Solution by HENRY HEATON, Belfield, N. D.

(1). The equation may be written $x/\sin x = 100$. As x varies from 0 to $\frac{1}{2}\pi$, $x/\sin x$ takes the successive values between 1 and $\frac{1}{2}\pi$. Hence the equation has no real root in the first quadrant. As x varies from $\frac{1}{2}\pi$ to π , $x/\sin x$ takes all the successive values from $\frac{1}{2}\pi$ to ∞ . Hence the equation has one real root in the second quadrant. In the third and fourth quadrants $x/\sin x$ is negative. Hence there can be no real root. At every round as x passes through the first and second quadrant $x/\sin x$ varies from ∞ to a minimum value then back to ∞ . This minimum value is in the first quadrant when $x = \tan x$. After the first round as long as $x < 100$ there are two real positive roots to every round. Hence there can be no real positive root when $x > 31\pi$, and the whole number of real positive roots is 31. The number of real negative roots is evidently the same with the same numerical values. Hence the whole number of real roots is 62. The largest real root evidently occurs between $x = 30\frac{1}{2}\pi$ and 31π . Put $x = y + 30\frac{1}{2}\pi$. Then $\sin(y + 30\frac{1}{2}\pi) = -\frac{1}{100}(y + 30\frac{1}{2}\pi)$.

$$\therefore \cos y = .958185 + \frac{1}{100}y.$$

If $\cos y = .95818$, $y = 16^\circ 38' = .2903^{(r)}$, a first approximate value of y . If $\cos y = .95818 + .002903 = .96108$, $y = 16^\circ 2' = .27983^{(r)}$, a second approximation.